

Induced wormholes due to quantum effects of spherically reduced matter in large N approximation

S. Nojiri*

*Department of Mathematics and Physics
National Defence Academy, Hashirimizu Yokosuka 239, JAPAN*

O. Obregon†

*Instituto de Fisica de la Universidad de Guanajuato
P.O. Box E-143, 37150 Leon Gto., Mexico*

S.D. Odintsov‡ and K.E. Osetrin§

*Tomsk State Pedagogical University
634041 Tomsk, RUSSIA*

Abstract

Using one-loop effective action in large N and s -wave approximation we discuss the possibility to induce primordial wormholes at the early Universe. An analytical solution is found for self-consistent primordial wormhole with constant radius. Numerical study gives the wormhole solution with increasing throat radius and increasing red-shift function. There is also some indication to the possibility of a topological phase transition.

*E-mail: nojiri@cc.nda.ac.jp

†E-mail: octavio@ifug3.ugto.mx

‡E-mail: odintsov@mail.tomsknet.ru

§E-mail: osetrin@tspu.edu.ru

It is a well known fact that spherical reduction of Einstein gravity (see, for example [1]) leads to specific 2d dilatonic gravity which belongs to the general class of 2d dilatonic gravities [2]. Spherical reduction of 4d matter leads to 2d dilaton coupled matter theories.

Recently, 2d conformal anomaly for dilaton coupled scalars has been found in ref.[3] (see also [4, 5, 6]). Integrating this conformal anomaly one finds the anomaly induced effective action [4, 5] (in s -wave and large N approximation) which was written in ref.[7] in most complete form. A wide spectrum of physical problems may be addressed using the above effective action. Let us only mention some of most interesting: anti-evaporation of multiple horizon black holes [7, 8]. Hawking radiation [9] in dilatonic supergravity [5], quantum cosmological models in dilatonic supergravity [10], study of semi-classical energy-momentum tensor in the presence of dilaton [11], etc.

In the present work we would like to discuss one more of these physical phenomena: the appearance of spherically symmetric wormholes due to the anomaly induced effective action. The wormholes represent handles of topological origin. They may be considered as bridges joining two different Universes or two separate regions of the same Universe [12]. It is known that wormholes cannot occur as solution of classical gravity-matter theory due to violations of energy conditions in classical relativity [12]. Hence, it is natural to expect their manifestations only at the quantum level (or yet in the theory like Born-Infeld D-brane or M-theory, for review, see [13]).

Recently, using quantum scalar stress-energy tensor calculated on spherically symmetric background in [14] the back reaction problem (i.e. self-consistent wormholes in semiclassical gravity) has been investigated in ref.[15] (see also [16]). It has been indeed numerically found a semiclassical quantum solution representing a wormhole connecting two asymptotically spatially flat regions [15]. This result is extremely interesting and it should be verified in another models and (or) approximations, taking into account the fact that the approach developed in [15] is not completely free from some serious drawbacks [16].

In the present work we show the possibility of inducing wormholes in the early Universe, making use of the effective action method [17] (large N and s -wave approximation is used).

We will start from the action of Einstein gravity with N minimal scalars

$$S = -\frac{1}{16\pi G} \int d^4x \sqrt{-g_{(4)}} \left(R^{(4)} - 2\Lambda \right) + \frac{1}{2} \sum_{i=1}^N \int d^4x \sqrt{-g_{(4)}} g_{(4)}^{\alpha\beta} \partial_\alpha \chi_i \partial_\beta \chi_i, \quad (1)$$

where χ_i are scalars, N is the number of scalars (in order to apply the large N approach, N is considered to be large, $N \gg 1$), G and Λ are the gravitational and cosmological constants.

The convenient choice for the spherically symmetric spacetime is the following one

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu + e^{-2\phi} d\Omega, \quad (2)$$

where $\mu, \nu = 0, 1$, $g_{\mu\nu}$ and ϕ depend only on x^0, x^1 and $d\Omega$ corresponds to the two-dimensional sphere.

The action (1), reduced according to (2) takes the form

$$S_{red} = \int d^2x \sqrt{-g} e^{-2\phi} \left[-\frac{1}{16\pi G} \{R + 2(\nabla\phi)^2 - 2\Lambda + 2e^{2\phi}\} + \frac{1}{2} \sum_{i=1}^N (\nabla\chi_i)^2 \right]. \quad (3)$$

Working in large N and s -wave approximation, one can calculate the quantum correction to S_{red} (effective action). Using 2d conformal anomaly for dilaton coupled scalar, calculated in [3] (see also [4, 5, 6]) one can find the anomaly induced effective action [4, 5] (with accuracy up to conformally invariant functional for the total effective action, see [17] for a review). There is no consistent approach to calculate this conformally invariant functional in closed form. However, one can find this functional as some expansion of Schwinger–DeWitt type [7] keeping only the leading term. Then, the effective action may be written in the following form [5, 7]¹

$$W = -\frac{N}{8\pi} \int d^2x \sqrt{-g} \left[\frac{1}{12} R \frac{1}{\Delta} R - \nabla^\lambda \phi \nabla_\lambda \phi \frac{1}{\Delta} R + \phi R + 2 \ln \mu^2 \nabla^\lambda \phi \nabla_\lambda \phi \right], \quad (4)$$

where Δ is two-dimensional laplacian, μ^2 is a dimensional parameter. Here, the first term represents the Polyakov anomaly induced action, the second and third terms give the dilaton dependent corrections to the anomaly induced action (these terms were found in ref.[5] and first discussed in connection with quantum gravity in ref. [6], similar terms with slightly different coefficients were derived in ref.[4]). The last term (conformally invariant functional) is found in ref.[7]. Note that it is possible to write the action (3) as some non-linear σ -model with one loop quantum correction (4). For this purpose, we rewrite (4) in a local form by introducing auxilliary fields u and v

$$W = -\frac{N}{8\pi} \int d^2x \sqrt{-g} \left[\frac{1}{12} R v - \nabla^\lambda \phi \nabla_\lambda \phi v + \phi R + 2 \ln \mu^2 \nabla^\lambda \phi \nabla_\lambda \phi + \nabla^\lambda u \nabla_\lambda v + u R \right]. \quad (5)$$

Then in the conformal gauge $g_{\mu\nu} = e^{2\rho} \bar{g}_{\mu\nu}$, we find

$$S_{red} + W = \int d^2x \sqrt{-\bar{g}} \left[\frac{1}{2} G_{ij}(X) \bar{g}^{\mu\nu} \partial_\mu X^i \partial_\nu X^j + \bar{R} \Phi(X) + T(X) \right], \quad (6)$$

where

$$\begin{aligned} X^i &= \{\phi, \rho, u, v, \chi_a\}, \\ \Phi(X) &= -\frac{1}{16\pi G} e^{-2\phi} - \frac{N}{8\pi} \left(\frac{v}{12} + \phi + v \right), \end{aligned}$$

¹Note that recently the effective action (4) has been rederived in ref.[18].

$$\begin{aligned}
T(X) &= -\frac{1}{16\pi G} \left(-2\Lambda e^{2\rho-2\phi} + e^{2\rho} \right), \\
G_{ij} &= \left(\begin{array}{cccc|c}
-\frac{e^{-2\phi}}{4\pi G} + \frac{N}{4\pi} (v - 2\ln \mu^2) & \frac{e^{-2\phi}}{4\pi G} - \frac{N}{4\pi} & 0 & 0 & 0 \\
\frac{e^{-2\phi}}{4\pi G} - \frac{N}{4\pi} & 0 & -\frac{N}{4\pi} & 0 & 0 \\
0 & -\frac{N}{4\pi} & 0 & -\frac{N}{8\pi} & 0 \\
0 & 0 & -\frac{N}{8\pi} & 0 & 0 \\
\hline
0 & 0 & 0 & 0 & e^{-2\phi}
\end{array} \right). \quad (7)
\end{aligned}$$

Working in the conformal gauge

$$g_{\pm\mp} = -\frac{1}{2}e^{2\rho}, \quad g_{\pm\pm} = 0, \quad (8)$$

the equations of motion may be obtained by the variation of $\Gamma = S_{red} + W$ with respect to $g^{\pm\pm}$, $g^{\pm\mp}$ and ϕ

$$\begin{aligned}
0 &= \frac{e^{-2\phi}}{4G} \left(2\partial_r \rho \partial_r \phi + (\partial_r \phi)^2 - \partial_r^2 \phi \right) \\
&\quad - \frac{N}{12} \left(\partial_r^2 \rho - (\partial_r \rho)^2 \right) - \frac{N}{2} \left(\rho + \frac{1}{2} \right) (\partial_r \phi)^2 \\
&\quad - \frac{N}{4} \left(2\partial_r \rho \partial_r \phi - \partial_r^2 \phi \right) - \frac{N}{4} \ln \mu^2 (\partial_r \phi)^2 + N t_0 \quad (9)
\end{aligned}$$

$$\begin{aligned}
0 &= \frac{e^{-2\phi}}{8G} \left(2\partial_r^2 \phi - 4(\partial_r \phi)^2 - 2\Lambda e^{2\rho} + 2e^{2\rho+2\phi} \right) \\
&\quad + \frac{N}{12} \partial_r^2 \rho + \frac{N}{4} (\partial_r \phi)^2 - \frac{N}{4} \partial_r^2 \phi \quad (10)
\end{aligned}$$

$$\begin{aligned}
0 &= -\frac{e^{-2\phi}}{4G} \left(-\partial_r^2 \phi + (\partial_r \phi)^2 + \partial_r^2 \rho + \Lambda e^{2\rho} \right) \\
&\quad + \frac{N}{4} \left\{ 2\partial_r (\rho \partial_r \phi) + \partial_r^2 \rho \right\} + \frac{N}{2} \ln \mu^2 \partial_r^2 \phi. \quad (11)
\end{aligned}$$

Here, t_0 is a constant which is determined by the initial conditions. Below we are interested in the static solution that is why we replace $\partial_{\pm} \rightarrow \pm \frac{1}{2} \partial_r$ where r is radial coordinate.

Combining (9) and (10) we get

$$\begin{aligned}
0 &= \frac{e^{-2\phi}}{4G} \left(-(\partial_r \phi)^2 + 2\partial_r \rho \partial_r \phi - \Lambda e^{2\rho} + e^{2\rho+2\phi} \right) \\
&\quad + \frac{N}{12} (\partial_r \rho)^2 - \frac{N}{2} \rho (\partial_r \phi)^2 - \frac{N}{2} \partial_r \rho \partial_r \phi - \frac{N}{4} \ln \mu^2 (\partial_r \phi)^2 + N t_0. \quad (12)
\end{aligned}$$

This equation may be used to determine t_0 from the initial condition, it decouples from the other two equations. Hence, Eq.(12) is not necessary in subsequent analysis.

Below, we consider a subclass of metrics (2) of the following type

$$ds^2 = -e^{2\rho} dt^2 + dl^2 + e^{-2\phi} d\Omega^2, \quad (13)$$

where $\rho = \rho(l)$, $\phi = \phi(l)$ and l is the proper distance. Note that to study exact solutions for matter in metrics of such sort, one can use the methods developed in ref.[19].

In order to come to metrics of the form (13), it is convenient to change the radial coordinate r to l by

$$dl = e^\rho dr . \quad (14)$$

Then we obtain

$$\partial_r = e^\rho \partial_l , \quad \partial_r^2 = e^{2\rho} (\partial_l^2 + \partial_l \rho \partial_l) . \quad (15)$$

Eqs. (10) and (11) may be rewritten as follows

$$\begin{aligned} 0 = & \left(\frac{e^{-2\phi}}{G} - N \right) \partial_l^2 \phi + \frac{N}{3} \partial_l^2 \rho + \left(-\frac{2e^{-2\phi}}{G} + N \right) (\partial_l \phi)^2 \\ & + \left(\frac{e^{-2\phi}}{G} - N \right) \partial_l \rho \partial_l \phi + \frac{1}{G} (-\Lambda e^{-2\phi} + 1) + \frac{N}{3} (\partial_l \rho)^2, \end{aligned} \quad (16)$$

$$\begin{aligned} 0 = & - \left(\frac{e^{-2\phi}}{G} - N \right) \partial_l^2 \rho + \left\{ \frac{e^{-2\phi}}{G} + N (2\rho + 2 \ln \mu^2) \right\} \partial_l^2 \phi - \frac{e^{-2\phi}}{G} (\partial_l \phi)^2 \\ & + \left\{ \frac{e^{-2\phi}}{G} + N (2\rho + 2 + 2 \ln \mu^2) \right\} \partial_l \rho \partial_l \phi \\ & - \left(\frac{e^{-2\phi}}{G} - N \right) (\partial_l \rho)^2 - \frac{e^{-2\phi}}{G} \Lambda . \end{aligned} \quad (17)$$

Our next task will be the study of Eqs. (16), (17) for the wormhole metric (13), where the usual notations are $f(l) = \exp(2\rho)$ (redshift function) and $r(l) = \exp(-\phi)$ (shape function). Actually, in our notations $r(l)$ is always positive, this function gives the wormhole throat.

First, we consider purely induced gravity, i.e. $N \rightarrow \infty$ case. Then Einstein action can be dropped away. For this case, the field equations have the form

$$0 = -\partial_l^2 \phi + \frac{1}{3} \partial_l^2 \rho + (\partial_l \phi)^2 - \partial_l \rho \partial_l \phi + \frac{1}{3} (\partial_l \rho)^2, \quad (18)$$

$$0 = \partial_l^2 \rho + (2\rho + a) \partial_l^2 \phi + (2\rho + 2 + a) \partial_l \rho \partial_l \phi + (\partial_l \rho)^2 . \quad (19)$$

where $a = 2 \ln \mu^2$. These equations admit the next integrals of motion

$$I_1 = e^\rho (\rho' + (2\rho + a)\phi') \quad (20)$$

$$I_2 = e^{2\rho} ((\rho')^2 - 3(2\rho + a)(\phi')^2 - 6\rho'\phi'), \quad (21)$$

here $' \equiv \partial_l$. In our study we take the following initial conditions:

$$f'(0) = r'(0) = 0 \quad (22)$$

For the conditions (22) we have $I_0 = I_1 = 0$, we get the trivial solution

$$\rho(l), \phi(l), \quad r(l), f(l) = \text{const.} \quad (23)$$

This indicates the possibility of inducing wormholes with finite throat.

For the case $f'_0 = r'_0 = 0$ from equations (16)–(17) in point $l = 0$ we have

$$0 = \left(\frac{e^{-2\phi_0}}{G} - N \right) \phi_0'' + \frac{N}{3} \rho_0'' + \frac{1}{G} (-\Lambda e^{-2\phi_0} + 1), \quad (24)$$

$$0 = - \left(\frac{e^{-2\phi_0}}{G} - N \right) \rho_0'' + \left\{ \frac{e^{-2\phi_0}}{G} + N(2\rho_0 + a) \right\} \phi_0'' - \frac{e^{-2\phi_0}}{G} \Lambda. \quad (25)$$

Let us consider the case when $\Lambda = 0$ and $r(l)$, $f(l)$ are non-decreasing functions near point $l = 0$, then we have the next restrictions for ρ_0 and ϕ_0

$$\begin{aligned} \phi_0 &\leq -\frac{1}{2} \ln(GN), \\ -\frac{e^{-2\phi_0}}{NG} - a - \frac{3}{N^2} \left(N - \frac{e^{-2\phi_0}}{G} \right)^2 &< 2\rho_0 \leq -\frac{e^{-2\phi_0}}{NG} - a. \end{aligned} \quad (26)$$

In other words, for the case $\Lambda = 0$ the throat radius is

$$r_0 \geq \sqrt{GN}. \quad (27)$$

Let us consider equations (16)–(17) for the case when $r(l) = r_0 = \text{const.}$ We get two solutions

1. $\Lambda \neq 0$, $|2GN\Lambda/3 - 2| \geq \sqrt{3}$,
 $r_0^2 = (2GN\Lambda/3 + 1 \pm \sqrt{(2GN\Lambda/3 - 2)^2 - 3}) / (2\Lambda)$,



$$f(l) = \begin{cases} (c_1 \cosh \sqrt{k} l + c_2 \sinh \sqrt{k} l)^2, & k > 0 \\ (c_1 + c_2 l)^2, & k = 0 \\ (c_1 \cos \sqrt{-k} l + c_2 \sin \sqrt{-k} l)^2, & k < 0, \text{ with} \end{cases}$$
 $k = (\Lambda r_0^2 - 1)/G$,
2. $\Lambda = 0$,
 $r_0 = \sqrt{GN}$, $f(l) = \left(\sqrt{f_0} \cos \sqrt{3/GN} l + (r_0 f'_0 / 2\sqrt{3f_0}) \sin \sqrt{3/GN} l \right)^2$.

Hence we found the wormhole (as a quantum effect induced object) with a constant throat radius and increasing or decreasing redshift function $f(l)$. This wormhole connects two spatial regions of spacetime. This can be viewed as a kind of infinitely long wormhole.

From another side in some above cases we get oscillating behaviour for $f(l)$. This fact indicates to some instability of wormhole configuration. Hence one can expect that kind of topological phase transition may occur and transform the wormhole to some another object, like black hole.

Let us discuss now the case when the classical gravity action is included, i.e. eqs.(16), (17) but r is not constant. In this case we perform a numerical study

of the equations of motion. Considering for simplicity $\Lambda = 0$ and $\ln \mu^2 = 1$, we present in figures 1. and 2. the typical result of our numerical study. The redshift function f increases as well as $r(l)$. It is interesting that qualitatively $r(l)$ and $f(l)$ behave in the same way as in ref.[15]. In particular, the redshift function does not approach to constant value at large l ., hence the whole metric is not asymptotically flat.



In these figures $r_0 = 27.1828$, $f_0 = 2.1405 \cdot 10^{-31}$, $r'_0 = f'_0 = 0$ and $N = 100$.

Hence, we showed that even for the whole system, i.e. classical gravity plus the quantum effective action due to matter, we can expect the appearance of primordial (induced) wormholes in the early Universe. The wormhole which appears

as result of quantum fluctuations connects two asymptotically (flat) regions of the early Universe. It may be with constant or increasing throat radius. The redshift function maybe increasing or oscillating. It would be very interesting to generalize above study for all types of quantum fields. Then one could investigate the connection between content of GUTs and inducing of quantum wormholes. For example, it may happen that SUSY GUTs better support inducing of wormholes as they include more fields then their non-SUSY versions.

Acknowledgments. SDO would like to thank S. Hawking and E. Mottola for useful discussion. O.O. was partially supported by a CONACYT Grant 28454-E. KEO was partially supported by a RFBR.

References

- [1] P. Hajicek, Phys.Rev. D30 (1984) 1178;
P. Thomi, B. Isaak and P. Hajicek, Phys.Rev. D30 (1984) 1168;
- [2] S.D. Odintsov and I.L. Shapiro, Phys.Lett. B263 (1991) 183;
T. Banks and M. O’Laughlin, Nucl.Phys.B362 (1991) 648;
- [3] E. Elizalde, S. Naftulin and S.D. Odintsov, Phys.Rev. D49 (1994) 2852;
- [4] R. Bousso and S. Hawking, Phys.Rev. D56 (1997) 7788;
- [5] S. Nojiri and S.D. Odintsov, Mod.Phys.Lett. 12 (1997) 2083; Phys.Rev. D57 (1998) 4847;
- [6] S. Nojiri and S.D. Odintsov, Phys.Rev. D57 (1998) 2363;
T. Chiba and M. Siino, Mod.Phys.Lett. A12 (1997) 709;
S. Ichinose, Phys.Rev. D57 (1998) 6224;
W. Kummer, H.Liebl and D.V. Vassilevich, Mod.Phys.Lett. A12 (1997) 2683;
J.S. Dowker, Class.Quant.Grav. 15 (1998) 1881;
- [7] S. Nojiri and S.D. Odintsov, hep-th/9806055, Phys.Rev.D to appear;
Mod.Phys.Lett. A13 (1998) 2695;
- [8] R. Bousso and S. Hawking, Phys.Rev. D57 (1998) 2436;
- [9] S. Hawking, Comm.Math.Phys. 43 (1975) 199;
- [10] S.J. Gates, T. Kadoyoshi, S. Nojiri and S.D. Odintsov, Phys.Rev. D58 (1998) 084026;

- [11] R. Balbinot and A. Fabbri, hep-th/9807123;
F.C. Lombardo, F.D. Mazzitelli and J.G.Russo, gr-qc/9808048;
- [12] M.S. Morris and K.S. Thorne, Am.J.Phys. 56 (1988) 395;
M.S. Thorne, K.S. Thorne and U. Yurtsever, Phys.Rev.Lett. 61 (1988) 1446.
- [13] G. Gibbons, preprint hep-th/9801106.
- [14] P. Anderson, W.A. Hiscock and D. Samuel, Phys.Rev. D51 (1995) 4337; B.E. Taylor, W.A. Hiscock and P. Anderson, gr-qc/9608036.
- [15] D. Hochberg, A. Popov and S.N. Sushkov, Phys.Rev.Lett. 78 (1997) 2050.
- [16] V. Khatsymovsky, In Proceedings of II Int. Conference Quantum Field Theory and Gravity, Tomsk, 1997; TGPU Publishing, Eds. I.L. Buchbinder and K.E. Osetrin, p.200.
- [17] I.L. Buchbinder, S.D. Odintsov and I.L. Shapiro, Effective Action in Quantum Gravity, IOP Publishing, Bristol and Philadelphia, 1992.
- [18] W. Kummer and D.V.Vassilevich, preprint hep-th/9811092;
- [19] V.V. Obukhov, Gravitation & Cosmology, 2(3) (1996) 256.